

(OT01) Discovery of ‘Z’
[25 marks]

You are given a sky map “Map-OT01” along with the question paper. This map shows only stars but no diffuse objects.

Four known stars (S1, S2, S3 and S4), which are present in the map above, are given in the table below with their Common names, Bayer designations and Equatorial coordinates.

[Σας δίνονται τα πιο κάτω 4 γνωστά άστρα με κοινινές σχετικά συντεταγμένες]

Sr. No.	Common Name	Bayer Name	RA	Dec
S1	Alpheratz	α Andromedae	00h 08m 24s	29° 05' 16"
S2	Markab	α Pegasi	23h 04m 46s	15° 12' 17"
S3	Scheat	β Pegasi	23h 03m 47s	28° 04' 58"
S4	Algenib	γ Pegasi	00h 13m 14s	15° 10' 59"

Complete the tasks (OT01.1) and (OT01.2) while you are planning the observations.

- (OT01.1) Your first task is to mark these 4 stars (with a circle around each star) and label them as S1, S2, S3 and S4 on the sky map “Map-OT01” provided to you. **[6]**

[Κυκλώστε και ονομάστε τα 4 άστρα S1-S4]

- (OT01.2) An astronomer has discovered a new diffuse object ‘Z’ at the following coordinates – **[7]**
RA: 21h 36m 10.6s, Dec: -26° 10' 24.4"
Mark the position of this diffuse object on the same sky map “Map-OT01” with a \oplus sign and label it as ‘Z’. Assume that a linear rectangular grid for equatorial coordinates is valid in the relevant region of the map.

[Εντοπίστε και σημειώστε στον χάρτη το αντικείμενο Z που σας δίνεται στις συγκεκριμένες συντεταγμένες]

The following task is to be performed once you reach the telescope station.

On the screen diagonally opposite to your station, initially a welcome message, followed by a sample sky (sky unrelated to the question) will be displayed along with a countdown timer. You may use this time to orient the telescope towards the screen and familiarize yourself with other equipment provided at the station. At the end of this time a part of the sky given in the map “Map-OT01” will be projected on the screen for the next 6 minutes. Note that the scale of the projection shown on the screen is different from the actual scale seen in the sky.



- (OT01.3) Find the new object ‘Z’ with the telescope using any appropriate eyepiece. Then centre the object properly in the field of view of the eyepiece with the crosshair, and show it to the examiner at your station. **[12]**
At the end of 6 minutes, the projection will be blurred for 20 seconds. At this point you must step away from the telescope. The projection will be restored for the examiner to check the view through the telescope. This marks the end of the first task.

[Στοχεύσετε στην απέναντι οθόνη, και εντοπίστε τον στόχο σας. Χρησιμοποιήστε μετά το μικρό προσοφθάλμιο για να φέρετε τον στόχο καλύτερα στο πεδίο όρασης]

(OT02) Lensing Time Delay

[25 marks]

Gravitational lensing can result in multiple images of a background source if the source, the lensing object and the observer are nearly aligned. These multiple images take different times to reach the observer, and if the background source is variable, each image shows the same feature in its variability after specific time delays. These time delay measurements are extremely useful to estimate the current expansion rate of the Universe, the Hubble constant.



We will consider the gravitational lens system shown in the above figure. The left hand panel shows a galaxy cluster (lens) together with 4 images of a background quasar formed due to gravitational lensing. The 4 images, labelled A, B, C and D, have different fluxes as each image is magnified by a different amount. For any given image the magnification does not change with time.

Light takes the largest time to travel for the image labelled D. The light coming from this quasar is variable, and astronomers have monitored this system for more than a decade now. The right hand panel of the figure shows the light curve for the image D.

On the screen opposite to your station you will see a movie of the gravitational lens system. This movie is 28 s long and will loop 6 times with a breaks of 1 minute or 2 minutes between runs. Every second on the watch corresponds to 250.0 days in the actual lens system.

[Λόγω του βαρυτικού φακού, οι πηγές A, B, C και D που πρόκειται για το ίδιο αντικείμενο, θα έχουν διαφορετικό χρόνο εκκίνησης της μεταβολής της φωτεινότητας τους. Είναι γνωστό πως η πηγή D θα είναι η τελευταία που θα αρχίσει να αλλάζει λαμπρότητα.]

- (OT02.1) Let the time delay of image D with respect to images A, B and C be given as $t_{DA} = t_D - t_A$, $t_{DB} = t_D - t_B$, and $t_{DC} = t_D - t_C$, respectively. Find these time delays taking any necessary steps to reduce the uncertainty in your results. [25]

[Καταγράψετε τον χρόνο έναρξης της μεταβολής στη φωτεινότητα της κάθε πηγής. Κατόπιν υπολογίστε την διάφορα χρόνου, την καθυστέρηση δηλαδή, από την πηγή D σε σχέση με τις άλλες 3 πηγές. Μετατρέψτε τα δευτερόλεπτα σε ημέρες.]